

# Degradation of the wetland sediment archive at Star Carr: An assessment of current palynological preservation

Bruce Alberta, Jim Innes, Jeff Blackford, Barry Taylor, Chantal Conneller, Nicky Milner

## Abstract

This paper presents the results of an investigation into the preservation status of pollen and other microfossils in the organic sediments at the wetland Mesolithic site of Star Carr. This study assesses the degradation of the pollen record in a profile at the edge of the archaeological site, adjacent to previous pollen work carried out from 1989 to 1991 and using it as a benchmark for comparison. There has been a severe degradation of pollen grains since the earlier work, with the upper peat devoid of pollen and the lower part of the organic profile badly affected. Only the very basal sediments retain well preserved pollen. Comparisons with hydrological and geo-chemical data obtained by other workers during the assessment of the Star Carr site suggest that oxidation caused by drainage and dessication of the organic sediments, perhaps originating in fissures in the drying peat, is a primary cause of the observed severe deterioration of the pollen record. Non-pollen palynomorphs (primarily fungal and algal spores) appear to be better preserved than pollen in the present bio-stratigraphic record, showing little surface degeneration, but are not recorded in the earlier work. The pollen archive in organic sediments at the Star Carr site is now badly damaged. Any further pollen work there should be undertaken urgently but is probably not justifiable.

## 1. Introduction

It is now six decades since the publication of Grahame Clark's monograph on excavations at the early Mesolithic archaeological site of Star Carr in the eastern Vale of Pickering in North-East England (Clark, 1954). It was the location of the Mesolithic settlement that first drew Clark to the site, as it lay on the edge of an Early Holocene lake (Candy et al., 2015, Palmer et al., 2015), now termed Lake Flixton, and so promised to provide the palaeoenvironmental data that he needed for his research on Mesolithic economy and land use. The iconic status enjoyed by Star Carr since Clark's work, therefore, is due not only to the prolific flint assemblages (Conneller and Schadla-Hall, 2003, Conneller et al., 2009), but also to the remarkable preservation and diversity of organic material associated with the site, which extended into the wetland sediments (Milner et al., 2011a). The organic components of Early Mesolithic material culture, which do not survive on dry sites, were found in abundance and in an excellent state of preservation, stratified within the waterlogged organic deposits that accumulated in the palaeolake margin and in the reedswamp, fen and carr wetland habitats associated with it. This stratification has allowed the use of palaeobotanical analyses, including both macrofossil (Dark, 2004, Taylor, 2011) and pollen (Walker and Godwin, 1954, Cloutman and Smith, 1988, Dark, 1998a, Dark, 1998b, Dark, 1998c, Cummins, 2000), to reconstruct the vegetation history around the site, providing an environmental context for the settlement at Star Carr (Innes et al., 2011, Milner et al., 2013). This combination of preserved organic cultural remains and multi-disciplinary study has made Star Carr the 'type' site for the British early Mesolithic, and a model for subsequent studies in wetland archaeology (Milner et al., 2011a). Not only could questions of site function, economy and seasonality of occupation be addressed, but the deeper understanding of the activities carried out at Star Carr allowed the site to be the hub of conceptual models of early Mesolithic land use, territoriality and interactions with the wider landscape (Clark, 1972).

### 1.1. Condition of the sediments

The continued existence of the palaeoenvironmental archive at Star Carr, and the potential for further multi-disciplinary analyses there, depends upon the quality of preservation of the waterlogged sediments and organic remains. Excavations in the mid 1980s (Schadla-Hall, 1987, Schadla-Hall and Cloutman, 1985, Mellars and Dark, 1998), however, showed that organic

preservation had deteriorated since Clark's work in 1950, and excavations since 2004 (Conneller, 2007, Milner et al., 2011b) have shown that the faunal and wood remains are now severely degraded. Drying, shrinkage and weathering of the peat has occurred and it is clear that modern land-use practices, particularly land drainage, have had an adverse effect on the hydrology and chemistry of the site and its environs (Brown et al., 2011), and therefore on the preservation of the organic material (Holden et al., 2006, Milner, 2007, Milner et al., 2011b). Installation of highly effective land drains in the last decade (Brown et al., 2011, Vorenhout, 2011) has accelerated this process, and the level of the peat surface has dropped considerably. Although the water level and the position of the edge of Lake Flixton fluctuated during the Early Mesolithic (Taylor, 2011), the edge remained a narrow zone and it is the deposits that formed in this lake-edge ecotone adjacent to the Mesolithic settlement (Dark, 1998a, Mellars, 1998) that today contain the archaeological organic remains. These shallow lake-edge peats are vulnerable to modern hydrological change, however, and are suffering the effects of drainage and falling watertable. The vulnerability of wetland archaeology to such damage is a problem at a national level, and many such sites and wetland landscapes are at risk (e.g. van de Noort et al., 2002, Brunning, 2013, Davis et al., 2015). At Star Carr, a national flagship site for wetland archaeology, the dessication of the organic sediments, and thus the degradation of the materials and information they contain, appears to have accelerated in the last decade, to the point where the archive of archaeological material is badly damaged (Milner et al., 2011b). Recent research at Star Carr, therefore, has concentrated on assessing the present condition of the organic sediments on and close to the archaeological site itself, to provide a benchmark to inform decisions regarding its future management and study (Emerick, 2011). Brown et al.'s (2011) work on the hydrology of the site and its catchment has shown that recent land drainage has been the cause of the dessication and oxidation of the peat causing chemical changes and promoting extreme sediment acidity. Boreham et al., 2011a, Boreham et al., 2011b performed a series of physical and geochemical analyses at the site that also showed severe acidification of the sediments due to lowered and fluctuating water tables. Oxidation and acidification have caused very serious and rapid decay of the antler, bone and wood remains (Milner et al., 2011a). In theory this very high acidity might not have such a severe effect upon pollen grain preservation, but watertable fluctuation, and associated peat dewatering, could well cause their corrosion (Lowe, 1982), and so the palynological record at Star Carr may be in as much danger as the rest of the organic material there. An assessment of current pollen preservation was therefore urgently required.

## 1.2. Previous palynological work

In this paper we present the results of an assessment of the current palynological status of the peat at Star Carr. We take as our benchmark the palynological analyses closest to the site itself (Day, 1993, Dark, 1998c) in the shallow lake margin peats adjacent to Clark's archaeological excavations (Fig. 1). Cloutman and Smith (1988) had earlier conducted pollen analyses on three profiles from a trench (VP85A) across the peat margin close to the site, and did not report any problems with poor pollen preservation. However, when Dark (1998c) performed more detailed pollen analyses from the same Trench (A), it was clear that the upper peats had been subject to drying and shrinkage. Dark (1998c) also noted a gradual deterioration of pollen and spore preservation with proximity to the surface, with well-preserved pollen being absent above 70 cm from the basal gravel. Dark (1998a) sampled at high resolution (approximately 2 to 4 year intervals) through the Mesolithic occupation phases, and identified phases of vegetation disturbance that coincided with increased microcharcoal percentages and also with the archaeologically-rich levels. Her interpretation that burning and significant disturbance of the local vegetation accompanied the phases of occupation at Star Carr was the first recognition of Early Mesolithic environmental impact there. During recent excavations in 2010 (Conneller et al., 2012) we recovered a peat profile from a new site, 20 m west of Dark's (1998a) profiles (Boreham et al., 2011b), and have conducted comparable analyses in order to investigate the condition of palynological remains, twenty years after Dark analysed her Star Carr samples.

## 2. Methods

A bulk sample of the basal 75 cm of sediment was recovered in aluminium monolith tins (numbered 321 and 322) from the southern end of excavation trench SC24 (Fig. 2), in a location selected to correspond closely to the altitude and position within the lake-edge peats of the Dark (1998a) profile. The surface altitude of the sampled location was 24.36 m OD. The peat between the top of the upper tin 322 and the ground surface was very dry and badly dessicated, and was not sampled. The sampled lower peat was from the edge of the archaeological site and contained occasional flint flakes, animal bone, and wood pieces; as such it was appropriate to sample in order to test the preservation of pollen and spores close to recent and future excavations. Its lithostratigraphy is shown in Table 1. Other monolith tins were collected from adjacent sections in case the first set proved barren of pollen, but these have not had to be used and are archived.

Pollen samples were analysed at 5 mm intervals, to provide a detailed assessment of microfossil preservation. Twelve samples are from the basal, minerogenic sediments and the remainder from the organic mud and peat above. Samples were prepared at the Department of Geography, University of Durham by means of a sequential application of cold HCL (10%), hot KOH (10%), hot HF (49%), warm HCL (10%) and a brief application of a hot acetolysis mixture (less than one-minute duration), the latter using a C<sub>4</sub>H<sub>6</sub>O<sub>3</sub> (90%) solution with H<sub>2</sub>SO<sub>4</sub> (10%). Pollen samples were filtered through a 175 µm sieve at the KOH stage, thus potentially reducing the number of charcoal particles above that size. After cleaning and dehydration (via ETOH), samples were mounted in high-viscosity silicone oil for light microscopic (LM) analysis. Both pollen and charcoal concentrations are calculated using an added exotic Lycopodium spike (Stockmarr, 1971). All Poaceae grains (including annular dimensions where grains are 34 µm or larger) as well as microcharcoal particles were measured along their maximum axis. The former procedure has been used to ascertain the relative importance of cereal-like (wild) grass taxa in the assemblage, and particularly *Glyceria fluitans* (cf. Albert and Innes, 2015), while measurements of microcharcoal are used to assess the relative proximity of fires, with high proportions of smaller charcoal elements indicating more distant fires (Blackford, 2000). Counts of 400 pollen grains were achieved in most cases. Non-pollen palynomorphs (NPPs), mainly fungal spores, have been identified (van Geel and Aptroot, 2006, Innes and Blackford, 2003). Most pollen and spore identification and nomenclature follows Moore et al. (1991) and was achieved at × 400 magnification on a standard light microscope, but higher magnifications were used to assess pollen preservation in more detail, according to degradation and damage. There has been considerable previous work on pollen preservation which has included a range of observational criteria, including folding, crumpling, breakage and surface corrosion (Cushing, 1967, Delcourt and Delcourt, 1980, Hall, 1981, Tipping, 1987, Twiddle and Bunting, 2010). These criteria have been used to characterise the mode of pollen deterioration - through transportation, in situ or progressive deterioration, and reworking and redeposition of grains. Studies have included species-specific (Campbell, 1999) and experimental approaches (Twiddle and Bunting, 2010). As the focus of this paper is the overall condition of the pollen archive, we have not presented deterioration data for individual taxa.

For this project, a specific aim has been to compare pollen preservation with the earlier study (Dark, 1998c), and assess the potential for future work. Hall (1981) concluded that the best indicators of deterioration were low pollen counts and high frequencies of deteriorated grains. In this study, comparison to the assemblages quantified by the earlier work at the same site can be added as the primary indicator of the deterioration since Dark's earlier samples were taken. In this study, degrees of crumpling, corrosion and folding of pollen are assessed using sub-samples of 50 grains, carefully examined at × 1000 magnification in immersion oil. Two classes of pollen degradation are recognised in this assessment, while relatively non-degraded grains are termed "Type 1" reflecting a well-preserved state in relation to each class (see Table 2). The first class of degradation is termed

“damage”, and includes mechanical folding (type 2) and tearing (Type 3) of grains, and may occur in situations involving movement or compression of sediment. The second class of degradation is termed “deterioration”, and reflects an erosion of fine micro-sculpturing elements of pollen exines (Type 2) or visible holes in the exine structure (Type 3) under  $\times 1000$  magnification. Such deterioration might result from either microbial (including fungal) agents or oxidation (Lebreton et al., 2010) of the exterior of the grain.

Counts of pollen, fungal remains and algal spores made at  $\times 400$  magnification were calculated in terms of percentage of total land pollen (TLP, c. 400 grains) and are shown in Fig. 3. Pollen condition assessment counts are based on the 50-grain sub-sample analysed at  $\times 1000$  magnification, shown as percentages. All pollen, spore and charcoal data have been plotted using the TILIA program (Grimm, 1993).

### 3. Results

The results of the pollen analysis are shown in Fig. 3. Six pollen zones are recognised based on changes in the main pollen taxa, with *Betula*, *Pinus*, *Salix* and *Poaceae* the major contributors, and these are similar to those of the earlier work by Dark (1998c), who recognised five major pollen zones. Her upper zone dominated by *Corylus* is missing from this new study at SC24, showing that the later pollen record that Dark recorded is no longer preserved. Most *Poaceae* grains in the SC24 profile are likely to derive from reedswamp grasses, and *Phragmites* remains are present in unit 2 of the lithostratigraphy (Table 1). A few grass grains can be identified as of *Glyceria* type (Albert and Innes, 2015). *Salix* values in the deepest zone from SC24 are higher than those of Dark (1998c) and may represent growth of *Salix* closer to the new sampling point, although the two sections are not far apart, as willow growth can be very localised. Minor taxa include many wetland herbs in low numbers, with *Cyperaceae* and *Filipendula* most abundant. These minor elements of the pollen assemblage are very similar to those identified by Dark (1998c). Results suggest a mosaic of wetland reedswamp, fen and carr vegetation with birch woodland on the dry land around the site. There are few pollen types present throughout the profile that indicate human impact on the environment, with only single grains of *Rumex acetosa* and *Plantago lanceolata*. As the new SC24 profile has no radiocarbon control, precise correlation with Dark's (1998c) pollen record is not possible but in general these new pollen results are very similar to her's, with Early Holocene (pre-Boreal) *Betula* woodland the local dominant vegetation outside the wetland. High *Filipendula* values near the base and the absence of a *Corylus* rise at the top of our diagram confirms that this assessment covers much of the first Holocene millennium, including the Mesolithic settlement phase, as does the dated diagram of Dark (1998b), and so comparisons can be made between the two data sets, although precise correlations are not possible. Microcharcoal results (Fig. 4) are also comparable, as might be expected from their proximity (Innes et al., 2004). Three microcharcoal peaks occur in the new profile, as in the previous work, interpreted as showing human activity at the site. Relatively little microcharcoal occurs in the upper part of the new sequence.

#### 3.1. Relative pollen degradation

In monolith 322 and the upper part of monolith 321, field observation had shown the peat to be severely dessicated (Fig. 2). At these depths, very little preserved pollen was found. Although similar depths above the peat base had been satisfactory for successful palynology in Dark's earlier profile, those sampled for the present study in 2010 were unable to yield a viable count. Only the lower 400 mm from tin 321, at depths of 725–1125 mm, contained pollen, although preservation levels are variable. Within this broad zone of some preservation, high levels of damaged grains have been recorded, with in some cases over 80% of the 50 grain subsample being damaged. The deteriorated category, most likely to represent post-depositional processes (Bunting and Tipping, 2000), includes many samples with over 60% deteriorated, and some over 80%. Whereas Dark (1998c) recorded <

5% deteriorated grains, high levels of deterioration and damage are present throughout this new profile, including some horizons without preservation. Total pollen concentration values in the section 725–1125 mm vary from sample to sample, with a range of 200,000 cm<sup>-3</sup> in basal layers to only 2000 in some upper levels. Comparable data from Dark (1998c) average 200,000, with peaks at 800,000 and minima around 50,000 (Dark, 1998c Fig. 11.3).

The pollen grains of *Pinus* species are considered more resistant than others to degradation (Sangster and Dale, 1961). The data in Fig. 3, Fig. 4 show higher levels of *Pinus* associated with low total concentrations of pollen, and higher percentages (60%+) of deteriorated grains, with *Pinus* levels sometimes in excess of 50% of TLP. Maximum values of 20% *Pinus* were reported by Dark (1998c) which has increased to 60% in SC24-e, suggesting that 50% of all other pollen has been lost, if *Pinus* pollen remains present in the same frequencies. *Juniperus*, a thin-walled grain that was present in Dark's (1998c) lower pollen zones, is entirely absent in SC24. In comparison to the original analyses of Dark (1998c), where only isolated crumpled/folded and deteriorated grains were noted, the present analyses produce a high encounter rate of moderately (Type 2) damaged and deteriorated grains. Moderately deteriorated pollen grains (with missing micro-sculpturing elements) are also common, averaging 20% of the sub-sample analysed, in basal levels below 1030 mm where degradation of pollen grains has been overall less severe.

In addition to the high levels of damage and deterioration (60%+) noted in those levels containing what appear to be inflated concentrations of *Pinus*, three distinct horizons lack countable levels of pollen, suggesting higher rates of degradation across all taxa. These horizons (bands 1–3 in Fig. 3, Fig. 4) are at 1015, 925–990 and 740–755 mm. Preservation was also very poor at 810 mm, although a low count was possible. Further evidence for degradation may be inferred from the unidentified fern spores (*Pteropsida* indet. cf. *Filicales*), which are considered resistant to oxidation and can be common in soil pollen analyses (Havinga, 1964, Tipping, 2000). The highest values (20–40% of TLP + taxon) are expressed at the boundaries of bands of very low concentration, where levels of severely degraded (Category 3) grains also increase. Dark (1998c) also recorded high values of *Pteropsida*, although these were in the upper levels of Trench A, M1 (top 400 mm of the profile), from a period classed here as non-polliniferous. In zones where Dark found abundant *Pteropsida* spores, this study shows uncountably low pollen frequencies, and in levels where Dark (1998c) found low background levels of generic ferns spores (< 2%) with occasional peaks, our samples recovered twenty years later contain 5–20%, as with the *Pinus* data suggesting loss of other taxa.

#### 4. Condition of the pollen archive

The lower 120 mm section of the sediment profile is miner-organic and has a moderately well-defined boundary to the overlying peat. Field estimation of water content while using the Troels-Smith scheme to record the lithostratigraphy (Table 1) during sample collection showed that these lowest levels were clearly saturated and levels of preservation of both macro- and microscopic material are good (section A on Fig. 2). Above these basal levels (section B on Fig. 2), however, handling of the peat showed that water content dropped sharply and although the peat was still damp, it was far less so than in section A. It also showed the peat in the upper tin 322 and in the top levels of tin 321 to be dry (section C on Fig. 2) and hardly damp at all to the touch. The peat from the top of tin 322 to the surface was extremely dry. Comparing the assessment profile with the descriptions and photographs of Dark's work and of the original excavations, it was clear that there has been a distinct fall in surface level, causing dessication and compaction of the sediments in this part of the site, presumably resulting from dewatering and shrinkage of the peat. This compaction has inflated the original concentrations and percentages of microcharcoal and other microfossils, exacerbated by the general pollen decay. Future microcharcoal results, either percentage or concentration, would therefore be skewed, not comparable to Dark's earlier data, and not an accurate record of original microcharcoal deposition. Our assessment analyses show that there are

major problems with pollen preservation in most parts of the sequence. Quality of preservation declines progressively higher in the profile, but not consistently, and there are groups of levels in tin 321 (bands 1, 2 and 3) where preservation was so poor that pollen was absent or at least uncountable. Significant levels of degradation or damage to pollen exines are, however, ubiquitous and are extremely high in several horizons within the profile. Sediments at the level of tin 322 are now virtually beyond use. Carbonate minerals reported by Dark in her earlier analyses were not found in our work, and the chemical changes noted by other studies (Boreham et al., 2011b) seem to have fundamentally altered the profile. French and Taylor (1985) have shown that the destructive effects of dehydration and oxidation begin immediately after sediments cease to be permanently waterlogged, even seasonally, so that even the lower levels at Star Carr are unlikely to sustain well-preserved pollen for very long. As Dark's (1998c) work did not include NPP analysis, no direct comparison can be made with the fungal spores recorded in the present study, which are therefore omitted from Fig. 4. It should be noted, however, that the fungal spores recorded throughout our counted levels show a much lower degree of surface degradation than the pollen, even in the upper parts of tin 321, and so fungal spores in general may well be robust and less susceptible to corrosion and deterioration. Further research is required on this subject.

#### 4.1. Causes of pollen degradation at Star Carr

The fact that recent pollen analyses of Star Carr peat sediments in Monolith 321 show major degradation of pollen grains merits some consideration as to agencies of preservation versus degradation. This can be done moreover in relation to hydrological (Brown et al., 2011) and geo-chemical (Boreham et al., 2011a, Boreham et al., 2011b) data so as to more strictly determine factors of a geo-chemical as opposed to biological degradation over the past twenty years. The relative importance of individual factors of degradation is not always easy to assess (Bryant and Holloway, 1983). Significantly, very low pH values determined to be pervasive in the case of Star Carr peats would indicate also a situation of very low biotic activity, and given a formerly near-pristine state determined in Dark's (1998c) analyses, a chemical agency during the last two decades is more likely. In the latter case, preservation of pollen by virtue of reduction in pre-drainage conditions in the valley is supported by regional hydrological data (Brown et al., 2011).

Regarding important patterning in the pollen preservation data (Fig. 4), that bands of poor preservation are present at various depths hints at factors of degradation beyond simple oxidation originating from surface layers, and might also result from oxidation originating in fissures in the peat bed that are sometimes observable in the profile as light sandy or silty clay layers (Table 1 and Fig. 2). Such layers form from expansion and contraction following dehydration and rehydration cycles, a property also observed in wood remains recently recovered from the Mesolithic site (Milner et al., 2011b). Oxidation in the Star Carr peat deposits is also measurable according to redox potential according to chemical analyses (Boreham et al., 2011b), including pH, iron oxide (esp. Iron II versus Iron III) content along multiple bore hole transects which lie between trench A and trench SC24. High redox potential values (Boreham et al., 2011b), but not pH per se although this influences redox (Boreham et al., 2011a), indicate severe oxidation, and most oxidized conditions are very positively correlated with total deterioration of pollen. Both mechanical damage and, more importantly, deterioration might be explained by a strong oxidizing reaction, given the extreme values. Redox (mV) values in excess of 400 (cf. western 5 m of Transect 2 near Monolith 321; Boreham et al., 2011a) appear to align with the pollen degradation Bands 2 and 3 reported in this study. Such a strong oxidizing reaction affecting the pollen is also enabled by the very high acidity of the sediment matrix, although this in itself is not an agent of degradation. It can be assumed that, because of the extremely acidic conditions that have developed at Star Carr during the past two decades, biotic agents of pollen degradation have been of lesser importance, as high acidity must have greatly reduced the rate of microbial activity in these sediments.

## 5. Conclusions

The assessment analyses in this study show that there are major problems with pollen preservation in many parts of the SC24 sequence, with only the base of the profile still maintaining a relatively close relationship to the more pristine spectra analysed earlier by Dark (1998c), although some less reliable analysis may still be practical in the lower 40 cm of the profile. Our sequence, located very close to Dark's original profile although not a duplicate section, shows clear evidence of substantial pollen deterioration since her work, to the extent that most parts of the profile are no longer viable for meaningful pollen analysis. While it is still possible to recover a pollen record that corresponds to the time of the Mesolithic occupation, the more fragile pollen types will have been differentially removed, preventing a reliable count, and all pollen types will have been affected by degradation. Although there may be pockets of better preservation, the section assessed in this study is likely to be representative of conditions at the wetland edge of the archaeological site as a whole, although preservation levels may well be better in deeper sediments further into the area of the palaeo-lake. Other more landward areas of the site, which appear even drier than the sampled location, may well be completely degraded. Preservation conditions appear to have been much better when Dark's analyses were completed and future deterioration is likely to continue unless preventative action is taken at the site. Nothing can be done about the damage that the pollen archive has already suffered, and any proposed palynological work around the Star Carr site itself should be done urgently, although our study indicates that degeneration has proceeded so far that it is already almost too late and any new pollen work would have to be recognised as compromised and unreliable. It is very likely that, nationally, the palaeoenvironmental resource at many other wetland archaeology sites, most of which have received less attention than Star Carr, are at similar risk and require urgent research attention before their palynological archive is destroyed.

## Acknowledgements

This work was funded by a NERC urgency grant (NE/I015191/1) and the Vale of Pickering Research Trust, University of Manchester, and University of York. We would like to thank the landowners of Star Carr, English Heritage and Natural England for granting permission to excavate. The figures were prepared in the Design and Imaging Unit, Geography Department, Durham University.

## References

- B.M. Albert, J.B. Innes Multi-profile, fine-resolution palynological and micro-charcoal analyses at Esklets, North York Moors, UK, with special reference to the Mesolithic-Neolithic transition. *Veget. Hist. Archaeobot.*, 24 (2015), pp. 357-375
- J.J. Blackford Charcoal fragments in surface samples following a fire and the implications for interpretation of subfossil charcoal data. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 164 (2000), pp. 33-42
- S. Boreham, J. Boreham, C.J. Rolfe Physical and chemical analyses of sediments from around Star Carr as indicators of preservation. *J. Wetl. Archeol.*, 11 (2011), pp. 20-35
- S. Boreham, C. Conneller, N. Milner, B. Taylor, A. Needham, J. Boreham, C.J. Rolfe. 2011b Geochemical indicators of preservation status and site deterioration at Star Carr. *J. Archaeol. Sci.*, 38 (2011), pp. 2833-2857
- A.G. Brown, C. Bradley, T. Grapes, I. Boomer. Hydrological assessment of Star Carr and the Hertford catchment, Yorkshire UK. *J. Wetl. Archeol.*, 11 (2011), pp. 36-55
- R. Brunning. Somerset's Peatland Archaeology: Managing and Investigating a Fragile Resource: The Results of the Monuments at Risk in Somerset Peatlands (MARISP) Project. Oxbow, Oxford (2013)
- V.M. Bryant, R.G. Holloway. The role of palynology in archaeology. *Adv. Archaeol. Meth. Theory*, 6 (1983), pp. 191-224

M.J. Bunting, R. Tipping. Sorting dross from data: possible indicators of post-depositional assemblage biasing in archaeological palynology. In, G. Bailey, R. Charles, N. Winder (Eds.), *Human Ecodynamics, Symposia of the Association for Environmental Archaeology No. 19*, Oxbow, Oxford (2000), pp. 63-68

I.D. Campbell. Quaternary pollen taphonomy: examples of differential redeposition and differential preservation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 149 (1999), pp. 245-256

I. Candy, A. Farry, C.M. Darvill, A. Palmer, S.P.E. Blockley, I.P. Matthews, A. MacLeod, L. Deepprose, N. Farley, R. Kearney, C. Conneller, B. Taylor, N. Milner. The evolution of Palaeolake Flixton and the environmental context of Star Carr: an oxygen and carbon isotopic record of environmental change for the early Holocene. *Proc. Geol. Assoc.*, 126 (2015), pp. 60-71

J.G.D. Clark. *Excavations at Star Carr*. Cambridge University Press, Cambridge (1954)

J.G.D. Clark. *Star Carr – a case study in bio-archaeology*. Addison Wesley Modular Publications No. 10, Cummings, Reading, Mass (1972)

E.W. Cloutman, A.G. Smith. Palaeoenvironments in the Vale of Pickering. Part 3: environmental history at Star Carr. *Proc. Prehist. Soc.*, 54 (1988), pp. 37-58

C. Conneller. New excavations at Star Carr. *PAST*, 56 (2007), pp. 3-5

C. Conneller, T. Schadla-Hall. Beyond Star Carr: the Vale of Pickering in the 10th millennium BP. *Proc. Prehist. Soc.*, 69 (2003), pp. 85-106

C. Conneller, N. Milner, T. Schadla-Hall, B. Taylor. The temporality of the Mesolithic landscape: new work at Star Carr. In. P. Crombé, M. Van Strydonck, J. Sergeant, M. Boudin, M. Bats (Eds.), *Chronology and Evolution Within the Mesolithic of North-West Europe, Proceedings of an International Meeting, Brussels, May 30th-June 1st 2007*, Cambridge Scholars Publishing, Newcastle upon Tyne (2009), pp. 77-94

C. Conneller, N. Milner, B. Taylor, M. Taylor. Substantial settlement in the European Early Mesolithic: new research at Star Carr. *Antiquity*, 86 (2012), pp. 1004-1020

G.E. Cummins. Fire! Accidental or strategic use of fire in the Early Mesolithic of the Eastern Vale of Pickering. In. *Mesolithic Lifeways. Current Research in Britain and Ireland*, Leicester Archaeology Monograph 7 (2000), pp. 75-84

E.J. Cushing. Evidence for differential pollen preservation in late quaternary sediments in Minnesota. *Rev. Palaeobot. Palynol.*, 4, The School of Archaeological Studies, University of Leicester (1967), pp. 87-110

P. Dark. Introduction and methods. In. P.A. Mellars, P. Dark (Eds.), *Star Carr in Context*, McDonald Institute for Archaeological Research, Cambridge (1998), pp. 111-117

P. Dark. Radiocarbon dating of the lake-edge deposits. In. P.A. Mellars, P. Dark (Eds.), *Star Carr in Context*, McDonald Institute for Archaeological Research, Cambridge (1998), pp. 119-124

P. Dark. Lake-edge sequences: results. In. P.A. Mellars, P. Dark (Eds.), *Star Carr in Context*, McDonald Institute for Archaeological Research, Cambridge (1998), pp. 125-146



- P. Dark. Plant remains as evidence for seasonality of site use in the Mesolithic period. *Environ. Archaeol.*, 9 (2004), pp. 39-45
- A. Davis, S. Bishop, J. Cheetham. Hydrological modelling of the Flag Fen archaeological site and wider landscape: main report. Final Report for Project 6187, Historic England, Swindon (2015)
- P. Day. Preliminary results of high-resolution palaeoecological analyses at Star Carr, Yorkshire. *Camb. Archaeol. J.*, 3 (1993), pp. 129-133
- P.A. Delcourt, H.R. Delcourt. Pollen preservation and quaternary environmental history in the southeastern United States. *Palynology*, 4 (1980), pp. 215-231
- K. Emerick. The management of Star Carr. *J. Wetl. Archeol.*, 11 (2011), pp. 120-132
- C. French, M. Taylor. Dessication and destruction: the immediate effects of dewatering at Etton, Cambridgeshire. *Ox. J. Archaeol.*, 4 (1985), pp. 139-155
- E.C. Grimm. TILIA Software. Illinois State Museum, Chicago (1993)
- Hall. Deteriorated pollen grains and the interpretation of quaternary pollen diagrams. *Rev. Palaeobot. Palynol.*, 32 (1981), pp. 193-206
- A.J. Havinga. Investigation into the differential corrosion susceptibility of pollen and spores. *Pollen Spores*, 6 (1964), pp. 621-635
- J. Holden, L.J. West, A.J. Howard, E. Maxfield, I. Panter, J. Oxley. Hydrological controls of in situ preservation of waterlogged archaeological deposits. *Earth Sci. Rev.*, 78 (2006), pp. 59-83
- J.B. Innes, J.J. Blackford. The ecology of late Mesolithic woodland disturbances: model testing with fungal spore assemblage data. *J. Archaeol. Sci.*, 30 (2003), pp. 185-194
- J.B. Innes, J.J. Blackford, I.G. Simmons. Testing the integrity of fine spatial resolution palaeoecological records: micro-charcoal data from near-duplicate peat profiles from the North York Moors, UK. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 214 (2004), pp. 295-307
- J.B. Innes, J.J. Blackford, I.G. Simmons. Mesolithic environments at Star Carr, the eastern Vale of Pickering and environs: local and regional contexts. *J. Wetl. Archeol.*, 11 (2011), pp. 85-108
- V. Lebreton, E. Messenger, L. Marquer, J. Renault-Miskovsky. A neotaphonomic experiment in pollen oxidation and its implications for archaeopalynology. *Rev. Palaeobot. Palynol.*, 162 (2010), pp. 29-38
- J.J. Lowe. Three Flandrian pollen profiles from the Teith valley, Perthshire, Scotland. II. Analyses of deteriorated pollen. *New Phytol.*, 90 (1982), pp. 371-385
- P.A. Mellars. Postscript: major issues in the interpretation of Star Carr. In: P.A. Mellars, P. Dark (Eds.), *Star Carr in Context*, McDonald Institute for Archaeological Research, Cambridge (1998), pp. 215-241
- P. Mellars, S.P. Dark. Star Carr in context: new archaeological and palaeoecological investigations at the Early Mesolithic Site of Star Carr, North Yorkshire. McDonald Institute Monographs, McDonald Institute for Archaeological Research, Cambridge (1998)
- N. Milner. Fading star. *Br. Archaeol.*, 96 (2007), pp. 10-14

- N. Milner, P. Lane, B. Taylor, C. Conneller, T. Schadla-Hall. Star Carr in a postglacial landscape: 60 years of research. *J. Wetl. Archeol.*, 11 (2011), pp. 1-19
- N. Milner, C. Conneller, B. Elliott, H. Koon, I. Panter, K. Penkman, B. Taylor, M. Taylor. From riches to rags: organic deterioration at Star Carr. *J. Archaeol. Sci.*, 38 (2011), pp. 2818-2832
- N. Milner, B. Taylor, C. Conneller, T. Schadla-Hall. *Star Carr: Life in Britain after the Ice Age*. Council for British Archaeology, York (2013)
- P.D. Moore, J.A. Webb, M.E. Collinson. *Pollen Analysis*. Blackwell, Oxford (1991)
- A.P. Palmer, I.P. Matthews, I. Candy, S.P.E. Blockley, A. Macleod, C.M. Darvill, N. Milner, C. Conneller, B. Taylor. The evolution of Palaeolake Flixton and the environmental context of Star Carr, NE. Yorkshire: stratigraphy and sedimentology of the Last Glacial-Interglacial Transition (LGIT) lacustrine sequences. *Proc. Geol. Assoc.*, 126 (2015), pp. 50-59
- A.G. Sangster, H.M. Dale. A preliminary study of differential pollen grain preservation. *Can. J. Bot.*, 39 (1961), pp. 35-43
- R.T. Schadla-Hall. Recent investigations of the early Mesolithic landscape in the Vale of Pickering. In: M. Zvevėbil, H. Blankholm (Eds.), *Mesolithic Northwest Europe: Recent Trends*, Department of Archaeology and Prehistory, University of Sheffield (1987), pp. 46-54
- R.T. Schadla-Hall, E.W. Cloutman. 'One cannot dig at random in a peat bog'. The eastern vale of Pickering and the archaeology of a buried landscape. M. Haselgrove, M. Millett, I.M. Smith (Eds.), *Archaeology From the Ploughsoil*, Department of Archaeology and Prehistory, University of Sheffield (1985), pp. 77-86
- J. Stockmarr. Tablets with spores used in absolute pollen analysis. *Pollen Spores*, 13 (1971), pp. 615-621
- B. Taylor. Early Mesolithic activity in the wetlands of the Lake Flixton basin. *J. Wetl. Archeol.*, 11 (2011), pp. 63-84
- R. Tipping. The origins of corroded pollen grains at five early postglacial sites in western Scotland. *Rev. Palaeobot. Palynol.*, 53 (1987), pp. 151-161
- R. Tipping. Pollen preservation analysis as a necessity in Holocene palynology. In: J.P. Huntley, S. Stallibrass (Eds.), *Taphonomy and Interpretation, Symposia of the Association for Environmental Archaeology 14*, Oxbow Books, Oxford (2000), pp. 23-33
- J. Troels-Smith. Characterisation of Unconsolidated Sediments. *Danm Geol Unders Series IV*, 3 (1955), pp. 38-73
- C.L. Twiddle, M.J. Bunting. Experimental investigations into the preservation of pollen grains: a pilot study of four pollen types. *Rev. Palaeobot. Palynol.*, 162 (2010), pp. 621-630
- R. van de Noort, W. Fletcher, G. Thomas, I. Carstairs, D. Patrick. *Monuments at Risk in England's Wetlands*. University of Exeter, Exeter (2002)
- B. van Geel, A. Aptroot. Fossil ascomycetes in quaternary deposits. *Nova Hedwigia*, 82 (2006), pp. 313-329

M. Vorenhout. In situ preservation and monitoring with particular application to Star Carr, Yorkshire, UK. *J. Wetl. Archeol.*, 11 (2011), pp. 56-62

D. Walker, H. Godwin. Lake stratigraphy, pollen-analysis and vegetational history. J.G.D. Clark (Ed.), *Excavations at Star Carr*, Cambridge University Press, Cambridge (1954), pp. 25-69